

## Distribution and ecological aspects of the herpetofauna of Strofadhés Islands (Ionian Archipelago, Greece)

Zur Verbreitung und Ökologie der Herpetofauna der Strophaden (Ionische Inseln, Griechenland)

EFSTRATIOS VALAKOS & MOYSIS MYLONAS

### ABSTRACT

The unknown herpetofauna of the Strofadhés Islands is presented and discussed. Its composition places Strofadhés Islands into the same biogeographical subunit as Peloponnesos and Zakynthos Island. The preferred habitats, densities, activities and thermal requirements for the most abundant species are given. The occurrence of gigantism in male individuals of *Podarcis taurica* in the smaller island (Arpya) is discussed.

### KURZFASSUNG

Die vorliegende Arbeit präsentiert Beobachtungen über die nahezu unbekannte Herpetofauna der Strophaden. Die Autoren diskutieren das Artenspektrum, welches die Strophaden als zur selben biogeographischen Unterinheit gehörig ausweist wie die Peloponnes und die Insel Zakynthos. Die Habitate, Populationsdichten, Aktivitäten und Temperatursprüche der häufigsten Arten werden beschrieben. Das Vorkommen von Rieswuchs bei männlichen *Podarcis taurica* wird diskutiert.

### KEYWORDS

Herpetofauna, Strofadhés Islands, Greece, distribution, ecology, gigantism; *Tarentola mauritanica fascicularis*, *Hemidactylus t. turcicus*, *Algyroides m. moreoticus*, *Podarcis taurica ionica*, *Telescopus f. fallax*.

### INTRODUCTION

On the contrary to the fair knowledge of the herpetofauna of the southern Ionian Islands (WERNER 1894, 1938; KEYMAR 1986) the herpetofauna of the Strofadhés is almost completely unknown. Only the species *Telescopus fallax* (FLEISCHMANN, 1831) was mentioned from Stamfani (WERNER 1938). The main reason for this ignorance which is extended also to the other animal and plant taxa is the island's distance from the nearest safe port

and the strong winds that usually blow in the area.

We visited the Strophadhés in December 1991 in the framework of a research program for the study of the fauna of small and remote Greek islands, supported by the "A. Leventis Foundation".

This work deals with the spatial distribution of the reptile species found and their ecology (preferred habitat, density, thermal ecology, activity).

### AREA AND METHODS

#### Area

The Strofadhés (Fig. 1) are the southernmost islands of the Ionian Archipelago, located 49 km south of Zakynthos Island and 51 km west of the Peloponnesos coasts. The bigger one of the two close

islands is named Stamfani (1.3 km<sup>2</sup>), and the smaller Arpya (0.2 km<sup>2</sup>). On Stamfani there is a monastery dating back to the 13th century. Today only one monk lives there and most of the cultivations and buildings have been ruined.

island. During the day the specimens were found under stones, in the crevices of stone walls and in the ruins. They were motionless with mean Tb 17.2°C while the corresponding Ta and Ts were 15.5°C and 15.7°C respectively. During the night we observed active individuals on the wall of the monastery. The mean Tb of the active animals was 15°C with corresponding Ta 14°C and Ts 14.8°C.

2. *Hemidactylus turcicus turcicus*  
(LINNAEUS, 1758)

Material: Four juvenile specimens (MS41291 19-22).

All individuals were found inactive under stones in the cultivated area around the monastery.

3. *Algyroides moreoticus moreoticus*  
BIBRON & BORY, 1883

The animals are ascribed to the subspecies *moreoticus* because of their white lateral ocelli (comp. KEYMAR 1986).

Material: Two adult males, two subadult females, one juvenile.

Sex	Inv. No.	SVL (mm)	L (mm)	W(g)
male	MS 412913	50.0	153.0	3.5
male	MS 412911	50.0	144.0	2.5
female	MS 412912	36.0	118.0	2.0
female	MS 412914	35.0	105.0	3.0
juvenile	MS 4129123	28.0	70.0	0.8

All three habitat types of the island are inhabited by this species.

The animals were active from 09.00 a. m. when Ta was 20°C till 04.00 p.m. The majority of the individuals was basking on stones close to the shrubs. Maximum activity was observed at 12.00 a.m. The Tb of the active animals was found to range from 20.2°C to 28°C (mean 24.8°C), whilst the corresponding Ta was 19°C - 21.2°C (mean 20.4°C), and Ts 19.2°C - 25°C (mean 21.3°C). Most of the active specimens were juvenile and subadult.

In a transect of 850 m long we ob-

served 22 *Algyroides*, four adults and 18 subadults and juveniles. The average density calculated was 5 specimens per 1000 m<sup>2</sup>.

4. *Telescopus fallax fallax*  
(FLEISCHMANN, 1837)

Material: One specimen (MS612913; 470 mm long, SQU 19, V 176, Sc 50/50, upper labials 8/8.

The animal was found inactive under a stone in the maquis.

B. Arpya Island

Two reptile species were found only: One gekkonid (*Tarentola mauritanica*), and the lacertid *Podarcis taurica*.

1. *Tarentola mauritanica fascicularis*  
(DAUDIN, 1802)

Material: Two adults and two juveniles.

Age	Inv. No.	SVL (mm)	L (mm)	W(g)
adult	SA 5129130	64.0	140.0	11.0
adult	SA 5129127	58.0	99.0	6.2
juvenile	SA 5129128	30.0	60.0	0.3
juvenile	SA 5129131	25.0	55.0	0.3

All specimens were found inactive, under stones, in the border zone of the island and around the abandoned cultivations. The Tb of a single specimen was found to be 17°C, corresponding Ta 14°C and Ts 15°C.

2. *Podarcis taurica ionica* LEHRS, 1902

Material: one adult male, two adult females, two subadults.

The lizards were found basking from 10.00 a.m. to 03.00 p.m. very close to the shrubs which offer them an effective shelter. No specimens were seen in the open field. The gigantic males (Fig. 2) were basking in the small openings between the shrub branches, never outside of the periphery of the shrubs.

Sex	Inv. No.	SVL (mm)	L (mm)	W(g)
male	SA 6129135	91.0	244.0	14.0
female	SA 5129126	52.0	130.0	2.3
female	SA 5129133	46.0	102.0	2.0
subad.	SA 5129132	41.0	131.0	1.5
subad.	SA 5129129	40.0	121.0	1.5

The mean Tb of the active animals was 25.6°C (24°C - 28°C), the cor-

responding mean of Ta was 20.6°C (16°C - 22°C) and of Ts 19.9°C (18°C - 21.8°C).

In a transect of 500 m long we observed 36 individuals. Four of them were gigantic males with total lengths of approximately 250 mm each. The estimated density of the population was 45 specimens per 1000 m<sup>2</sup>.

## DISCUSSION

The great depths that surround the Strofadhes and exceed 900 m in the direction of Peloponnesos and Zakynthos, the Tyrrhenian limestone that constitutes the geological formation of the islands' surface (PSARIANOS & al. 1979), and the shallow sea, not deeper than 8 m, that separates the islands, lead us to conclude that:

The Strofadhes are quite recent islands, not older than 350.000 years as the surfaces of the islands consist of Tyrrhenian sediments. The islands were joined together at least during Würm glaciation, 15.000 years before present, but were never connected to the Peloponnesos or Zakynthos.

Both fauna and flora are expected to originate rather recently from adjacent land species that managed to pass the existing sea barrier.

Consequently, endemism - especially above species level - is expected to be less pronounced than in other Greek islands.

The herpetofauna observed on the Strofadhes verifies the above expectations:

All species found are also inhabitants of the adjacent coasts of the Peloponnesos and Zakynthos; especially *Algyroides m. moreoticus* is a characteristic endemic species in this area.

The distribution of the Lacertidae on the two islands of Strofadhes (*A. m. moreoticus* exclusively in Stamfani, *P. taurica ionica* in Arpya only) is of special interest. It can be attributed either to biogeographical (dispersal) or ecological (available niche, competition, enemies) reasons. ARNOLD (1987) and BUTTLE (1988) refer to many cases of real sympat-

try of these two species in Peloponnesos without any indication of mutual ecological replacement. The ecological conditions on both islands (weather, soil, vegetation type) are more or less similar, except for the open cultivated areas on Stamfani. For all that, *P. taurica* which usually prefers the open grassland (CHONDROPOULOS 1984) lives on Arpya where no grass exists, whilst is absent from Stamfani where the cultivations would constitute a suitable habitat. As both islands are very close to each other and must have been united until quite recently, the reasons for the specific separation cannot be biogeographical. Furthermore it seems improbable that a species can disperse 50 km across the sea - passively in this case - and, on the other hand, should be incapable to cross 800 m between the islands. Such a problematic distribution has been observed in the Balearic Islands concerning *P. lilfordi* (GÜNTHER, 1874) and *P. sicula* RAFINESQUE, 1810 (PEREZ MEL-LADO 1989). We believe that these distributional questions could be answered by special ecological research suited for very small islands and islets, not by standard ecological approaches.

The observed activities of *A. moreoticus* and *P. taurica* are within the frame of the known winter activities of Lacertidae in South Europe (AVERY 1982; VALAKOS 1990). The animals are active during the warm hours in the middle of the day, having lower body temperatures than in other periods of the year, when they are basking late in the morning and early in the afternoon.

The density of *P. taurica* on Arpya is almost twice as high as the known densities of this species in other Ionian Islands and Western Greece as are given by CHONDROPOULOS (1984). No comparative data exist for *A. moreoticus*.

*T. mauritanica* is a typical nocturnal species. It is inactive during the day, even though its body temperature is higher than (17.2°C) than at night (15°C), when the animal is active. The same has been reported by ARNOLD & BURTON (1978)

for other populations.

The adult males of *P. taurica ionica* in Arpya can be characterized gigantic, compared to males of other populations of the subspecies. In table 1 we give L, SVL, and W of the biggest males observed in different populations of *P. taurica ionica*. In specimens of the western Peloponnesos there is also an increase in body length, but not correspondingly in body weight as in Arpya.

Table 1: Dimensions of the biggest known males of *Podarcis taurica ionica* (LEHRS, 1902) from different regions.

Tabelle 1: Maße der größten bekannten Männchen von *Podarcis taurica ionica* (LEHRS, 1902) aus verschiedenen Gebieten.

Region	L (mm)	SVL (mm)	W (g)	Reference
Zakynthos Isl.	232.5	73.6	7.6	CHONDROPOULOS (1984)
Kefalonia Isl.	232.5	75.5	9.2	CHONDROPOULOS (1984)
Ithaki Isl.	217.5	70.2	7.5	CHONDROPOULOS (1984)
Kefalonia Isl.	232.5	75.5	9.2	CHONDROPOULOS (1984)
Kerkyra Isl.	213.5	71.9	6.7	CHONDROPOULOS (1984)
W. Peloponnesos	241.5	84.9	10.8	CHONDROPOULOS (1984)
Epirus	220.9	73.5	7.6	CHONDROPOULOS (1984)
Arpya Isl.	244.0	91.0	14.0	this work




Fig. 2: a - The gigantic male of *Podarcis taurica ionica* (LEHRS, 1902) collected in Arpya; b - Male of *P. taurica ionica* from NW-Peloponnesos.

Abb. 2: a - Das riesenwüchsige Männchen von *Podarcis taurica ionica* (LEHRS, 1902) von der Insel Arpya; b - Männliche *P. taurica ionica* von der NW-Peloponnes.

The problem of differentiation of the body size with insularity has been studied by many scientists (e. g. SACCHI 1961; ANGERBJORN 1986). RADOVANOVIC (1956) refers to island populations of gigantic *P. sicula* in Dalmatia.

We consider that gigantism in the studied case could be attributed to the ab-

sence of predation and interspecific competition as documented by ANGERBJORN (1986) for wood mice (*Apodemus*), or to the increasing intraspecific competition because of increasing density related to the small size of the islands, as reported by MELTON (1982).

#### ACKNOWLEDGEMENTS

We are grateful to the "Leventis Foundation" for the economical support, and to the colleagues of

the scientific expedition to the Strofadhés Islands for their collaboration.

#### REFERENCES

- ANGERBJORN, A. (1986): Gigantism in island populations of the wood mice (*Apodemus*) in Europe.- *Oikos*; 47:47-56.
- ARNOLD, E. N. (1987): Resource partition among lacertid lizards in southern Europe.- *J. Zoology*, London; (B) 1:739-782.
- ARNOLD, E. N. & BURTON, J. A. (1978): A field guide to the Reptiles and Amphibians of Britain and Europe; London (Collins), 272 pp.
- AVERY, R. A. (1982): Field studies of body temperatures and thermoregulation. In: GANS, C. & POUGH, F. (Eds.): *Biology of the reptilia* Vol. 13 D; London, New York (Academic Press), pp.93-166.
- BUTTLE, D. (1988): Further notes on reptiles and amphibians of the Peloponnese.- *British Herpetol. Soc. Bull.*; 26:14-20.
- CHONDROPOULOS, P. V. (1984): Study on the lizards Lacertidae in West Greece; Ph D. Thesis (Univ. Patras), 196 pp.
- KEYMAR, P. F. (1986): Die Amphibien und Reptilien der Ionischen Region (Griechenland). Analyse ihrer rezenten Verbreitungsmuster und Überlegungen zu ihrer Ausbreitungsgeschichte.- *OGH-Nachrichten*, Wien; (8/9):8-44.
- KREBS, C. J. (1987): *Ecological methodology*; New York (Harper & Row), 652 pp.
- MELTON, R. H. (1982): Body size in island *Peromyscus*: A pattern and a hypothesis.- *Evolutionary Theory*; 6:113-126.
- PEREZ MELLADO, V. (1989): Estudio ecológico de la lagartija Balear *Podarcis lilfordi* (GÜNTHER, 1874) en Menorca.- *Revista de Menorca*; 80:455-511.
- PSARIANOS, P. & VERGINIS, S. & VERYKIOU, E. & LIVADITIS, G. (1979): Geographie und Geologie der Inseln Strophades.- *Ann. Géologiques des Pays Helléniques*; 29:729-739.
- RADOVANOVIC, M. (1956): Zum Problem der Speziation bei Inseleidechsen.- *Zool. Jahrb.*; 86:395-436.
- SACCHI, C. F. (1961): Considérations sur les phénomènes microévolutifs d'animaux des petites îles Méditerranéennes.- *Colloques internationaux du C.N.R.S. (Les peuplement des îles Méditerranéennes)*, Banyuls-Sur-Mer; 94:321-337.
- VALAKOS, E. (1990): The ecology of the lizard *Podarcis erhardii* (BEDRIAGA, 1882) (Sauria: Lacertidae) in a typical insular ecosystem on Naxos Island (Greece); Ph. D. Thesis (Univ. Athens), 214 pp.
- WERNER, F. (1894): Die Reptilien- und Batrachierfauna der jonischen Inseln.- *Sitz.-ber. Zool. Bot. Ges. Wien*; 44:225-237.
- WERNER, F. (1938): Die Amphibien und Reptilien Griechenlands.- *Zoologica*, Stuttgart; 35(94):1-117.

DATE OF SUBMISSION: January 10th, 1992

AUTHORS: Dr. Efstratios VALAKOS & Dr. Moysis MYLONAS, University of Athens, Biological Department, Section of Ecology and Taxonomy, Panepistimioupolis, Ilissia 157-71, Athens, Greece.